# **DuPont™** RelyOn™ Multi-Purpose Disinfectant Cleaner

BROAD SPECTRUM DISINFECTANT

Virucidal, Bactericidal, Germicidal, Fungicidal Compliant with OSHA bloodbourne pathogen standard Cleans and disinfects in one step Compatible with most non-porous surfaces NSF I isted

#### **ACTIVE INGREDIENTS**

Potassium peroxymonosultate	21.41%
Sodium chloride	1.50%
INERT INGREDIENTS	77.09%
TOTAL	.100.00%
Equivalent to 9.75% Available Chinrine	

### KEEP OUT OF REACH OF CHILDREN

## DANGER/PELIGRO

TABLET FORM

#### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for further treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for further treatment advice.

IF SWALLOWED: Call Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Powder is corrosive. Causes skin burns and irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear protective clothing and rubber gloves. Wear goggles, face shield or safety glasses. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Corrosive statement does not refer to 1% in-use solution.

### STORAGE AND DISPOSAL

Store in a cool dry place in tightly closed container. Keep out of reach of children. Always replace lid after use. Retain dessicant cannister with product during storage. Wash empty container thoroubly and dispose of in trash. Do not mix this product with other chemicals.

#### FOR USE IN-

Cleaning and disinfecting hard, non-porous surfaces in institutional, commercial, residential and public service facilities including hospitals and nursing homes; field medical units, medical and dental offices; clinics and other health-care facilities; Emergency Medical Services (EMS) facilities, equipment and gear including respirators; laboratories; food processing plants, equipment and warehouses; food service and preparation areas including commercial and institutional kitchens, restaurants and cafeterias; restrooms and portable restrooms; retail stores including supermarkets, florists, barbers and beauty salons; factories; schools and daycare centers; hotels; offices; correctional facilities; funeral parlors and mortuaries; gymnasiums, stadiums and health spas; transportation terminals and vehicles including ships, automobiles, trucks and trains, and homes and residences.

# EFFECTIVE AGAINST A WIDE RANGE OF VIRUSES\*, BACTERIA AND FUNGI:

\* Hepatitis A Virus (HAV), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), and HIV (Human Immunodeficieny Virus (HIV-1) (AIDS Virus) Influenza A Virus, Respiratory Syncytial Virus (RSV), Norovirus (Norwalk-like Viruses), Avian Influenza A Virus

Bacillus cereeus, Campylobacter jejuni, Chlamydia psittaci, Clostridium perfringens, Listeria monocytogenes, Shigella sonnei, Streptococcus pyogenes, Helicobacter pylori, Klebsiella pneumoniae, Escherichia coli 0157:H7, Escherichia coli, Salmonella typhimurium, Salmonella choleraesuis, Pseudomonas aeruginosa, Staphylococcus aureaus Staphylococcus aureaus (MRSA), Vancomycin-Resistant Enterococcus faecalis (VRE), Candida albicans, and Trichophyton mentagrophytes

#### DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

# INSTITUTIONAL, FOOD PROCESSING, AND PUBLIC SERVICE FACILITIES:

With DuPont™ RelyOn™ Multi-Purpose Disinfectant Cleaner only one product is needed to clean and dikinfect.

Do Not Use DuPont™ RelyOn™ Multi-Purpose Disinfectant Cleaner on acid-sensitive surfaces such as marble or soft metals such as copper, brass, or certain grades of aluminum. Avoid splashing DuPant™ RelyOn™ Multi-Purpose Disinfectant Cleaner solution on textiles or carpets. End user should assure compatibility of all critical materials that may be contacted by DuPont™ RelyOn™ Multi Purpose Disinfectant Cleaner. Cleaning Procedure: Remove gross dirt and prepare a 1.0% solution according to the table below. Apply to surface using a sponge, brush, or spray device, scrubbing as necessary until the surface is visibly clean. Rinse.

CLEANING AND DISINFECTING HARD NON-POROUS FOOD CONTACT SURFACES: Remove gross dirt and prepare 1.0% solution according to the table below. Apply to surface using a sponge, brush, or spray device until the surface is visibly clean. Let stand for 10 minutes and rinse with potable water.

CLEANING AND DISINFECTING HARD NON-POROUS NON-FOOD CONTACT SURFACES: Remove gross dirt and use 1.0% DuPont<sup>TM</sup> RelyOn<sup>TM</sup> Multi-Purpose Disinfectant Cleaner solution prepared according to the table below. Apply to surface using a mop, sponge, brushes or spray device until the surface is visibly clean. Let stand 10 minutes and air dry. In cases of fungal contamination of non-food contact surfaces, follow these instructions substituting a 2.0% DuPont<sup>TM</sup> RelyOn<sup>TM</sup> Multi-Purpose Disinfectant Cleaner Solution.

This product is not to be used on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

CLEANING AND DISINFECTING MANIKINS USED IN CPR TRAINING: Sanitizing manikins between students.

Prepare a 3% solution of DuPont™ RelyOn™ Multi-Purpose
Disinfectant Cleaner (according to dilution chart) by dissolving 3 tablets (5gm each) of DuPont™ RelyOn™ Multi-Purpose
Disinfectant Cleaner in one pint (16 02) water or by dissolving 1 tablet (5 gm) of DuPont™ RelyOn™ Multi-purpose Disinfectant cleaner in premarked 5.3 oz spray bottle that has been filled to the line with water. Thoroughly spray the manikin surface (mouth and nose area) or wipe with a cloth saturated with 3% solution until fully wetted. Allow surface to remain visibly wet for a minimum of 30 seconds, and wipe with a disposable cloth until dry. Rinse

CRISIS SCENE DISINFECTION OF HARD NON-POROUS SURFACES: For bacterial and viral\* conrol: Streptococcus pyogenes, Helicobacter pylori, Klebsiella pneumoniae, Escherichia coli Salmonella typhimurium, aureus, Methicillin-Resistant Staphylococcus aureus (MRSA), Vancomycin-Resistant Enterococcus faecalis, (VRE), and Staphylococcus epidermidis. \*Hepatitis A Virus (HAV), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), and HIV (Human Immunodeficiency Virus (HIV-1) (AIDS virus), Influenza A Virus, Respiratory Syncytial Virus (RSV), Norwalk-like viruses), Avian Influenza A Virus.

surface with potable water.

For a 1% solution from tablet: Remove gross dirt. Add one 5gm (0.18 oz) DuPont™ RelyOn™ Multi-Purpose Disinfectant Cleaner tablet per pint of water. Cap container and gently shake until tablet is dissolved. Apply solution to surface using a sponge, brush, or. spray device until the surface is visibly wet. Let stand for 10 minutes and air dry.

# SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST \*HIV-1, HEPATITIS B (HBV), AND HEPATITIS C (HCV) ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

\*Kills HIV-1, HAV, HBV, and HCV on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings (Hospitals, Nursing Homes, etc) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/ objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects using a cleaner (such as DuPont™ RelyOn™ Muti-purpose Disinfectant Cleaner) before application of this product as a disinfectant (see previous label instructions for use of DuPont™ RelyOn™ Multi-purpose Disinfectant Cleaner as cleaner)

CONTACT TIME: Allow surface to remain wet for 10 minutes.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

10 Tbsp

10 Tbsp

2-1/2 cups

2-1/2 cups

Net Contents: refer to container label

Choose a delivery opti	on from the list below	Add the following amount of DuPont™ RelyOn™ Multi-Purpose Disinfectant Cleaner to the designated volume of water									
		1%									
DuPont™ RelyOn™ M	ulti-Purpose Disinfectant Cleaner	16 oz. (Pint)	32 oz. (Quat)		128 oz. (Gallon)		160 oz. (5 Quarts)		640	640 oz. (5 Gallons)	
Metric	Standard										
5 g Tablet	0.18 oz Tablet	1 Tablet 2 Tablets 8		8 Tablets 10		10 Tab	0 Tablets 4		40 Tablets		
40 g Tablet	1.44 oz Tablet		1 Tablet		5 Tablets		blets				
50 g Sachet	g Sachet 1.8 oz Sachet						1 Sachet		4 Sachets		
500 g Shaker Bottle	17.6 oz Shaker Bottle	1 tsp 2 tsp		8 tsp		10 tsp		401	40 tsp		
5 kg Bucket	176 oz Bucket	1 tsp	2 tsp		8 tsp		10 tsp		40 tsp		
		2%									
5 g Tablet	ablet 0.18 oz Tablet		4 Tablets		16 Tablets		20 Tablets		80	80 Tablets	
40 g Tablet	1.44 oz Tablet			2 Tablets		<u>-</u>			10 Tablets		
50 g Sachet	1.8 oz Sachet				Ì	2 Sach		iets 8		achets	
500 g Shaker Bottle	17.6 oz Shaker Bottle	Bottle 2 ts 4 ts		4 tsp 16 tsp (1/3 cu		cup) 20 tsp			1-2/3 cups		
5 kg Bucket	176 oz Bucket	2 tsp	4 tsp		16 tsp (1/3 cup)		20 tsp		1-2/3 cups		
		3%									
		170 ml (5.75 oz)	16 oz. (Pint)	32 oz.	(Quart)	128 oz. (Ga	llon)	160 oz. (5 Quart	s)	640 oz. (5 Gallon)	
5 g Tablet	0.18 oz Tablet	1 Tablet	3 Tablets	6 Tab	ets	24 Tablets		30 Tablets		120 Tablets	
40 g Tablet	40 g Tablet	l				3 Tablets		<u> </u>		15 Tablets	
50 g Sachet	50 g Sachet							3 Sachets		12 Sachets	

2 Tbsp

2 Tbsp

0.5 cups

0.5 cups

1 Tbsp

1 Tbsp

E.I. du Pont de Nemours and Company 1007 Market Street Wilmington, DE 19898

Questions? Call 1-800-441-7515

Manufactured for:

EPA Reg. No.71654-7 EPA Est. No. 62432-EN-001 tab insert 072706 Rev 12/6/06 A00964807/RelyOnT/IFU/Dupont,USA/12.06/P

tsp = teaspoon Tbsp = tablespoon

5 kg Bucket

500 g Shaker Bottle

500 g Shaker Bottle

5 kg Bucket